



INDICATIONS FOR USE

To prevent reinfection of *Toxocara canis* in puppies and adult dogs and in lactating bitches after whelping. For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal examination. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

ACTIVE INGREDIENT: 2.27 mg of pyrantel pamoate base as pyrantel pamoate per mL.



Manufactured for
Farnam Pet Products
A Division of Farnam Companies, Inc.
P.O. Box 34820
Phoenix, AZ 85067-4820
For information or comments, call
1-800-234-2269
Visit our Web site at famampet.com

Product Code: 3099 500021 N-Rev0707
©2007 Farnam Companies, Inc. 07-1454

NDC 017135-703-12

D-Worm™ 60
(pyrantel pamoate)

Hookworms & Large Roundworms
Liquid
for Puppies & Dogs



Dog Dewormer

Easy-to-use

- Removes **Hookworms and Large Roundworms (Ascarids)**
- Prevents reinfection from **Roundworms**

5 ml per 5 lb Body Weight

ANADA #200-248 • Approved by FDA
NET CONTENTS: 60 mL (2 fl oz)



CAUTION:
IT IS EXTREMELY IMPORTANT
THAT THE PRODUCT BE SHAKEN THOROUGHLY
BEFORE EVERY USE.

DIRECTIONS FOR USE

For maximum control and prevention of reinfection, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfection. Administer one full teaspoonful (5 mL) for each 5 lb of body weight.

For the removal of large roundworms (ascarids) and hookworms. Administer one full teaspoonful (5 mL) for each 5 lb of body weight. It is not necessary to withhold food from your dog prior to treatment. Dogs usually find this wormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption.

It is recommended that dogs maintained under conditions of constant exposure to worm infection should have a follow-up fecal exam within 2 to 4 weeks after first treatment.

IMPORTANT: SHAKE WELL BEFORE USE
READ ACCOMPANYING PACKAGE INSERT

FOR ANIMAL USE ONLY
RECOMMENDED STORAGE:
Store Below 86° F (30° C).

WARNING: KEEP OUT OF REACH OF CHILDREN

NDC 017135-703-12

D-Worm™ 60
(pyrantel pamoate)

Hookworms & Large Roundworms
Liquid
for Puppies & Dogs



Dog Dewormer

Easy-to-use

- Removes **Hookworms and Large Roundworms (Ascarids)**
- Prevents reinfection from **Roundworms**

5 ml per 5 lb Body Weight

ANADA #200-248 • Approved by FDA
NET CONTENTS: 60 mL (2 fl oz)



Hookworms & Large Roundworms
Liquid Dewormer for Puppies & Dogs

5 ml per 5 lb Body Weight

Hookworms & Large Roundworms
Liquid Dewormer for Puppies & Dogs

5 ml per 5 lb Body Weight




INDICATIONS FOR USE: To prevent reinfection of *Toxocara canis* in puppies and adult dogs and in lactating bitches after whelping, for the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Unteria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by bloody fecal examination. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

IMPORTANT: SHAKE WELL BEFORE USE
READ ACCOMPANYING PACKAGE INSERT
FOR ANIMAL USE ONLY

RECOMMENDED STORAGE:
 Store Below 86°F (30°C).
WARNING-KEEP OUT OF REACH OF CHILDREN

Product Code: 3009
 Lot No. _____
 Exp. Date _____
 ©2007 Farnam Companies, Inc.
 07-1560



D-Worm™ 60
(pyrantel pamoate)
Dog Dewormer
 Liquid for Puppies & Dogs

Easy to Use

- Removes Hookworms and Large Roundworms (Ascarids)
- Prevents reinfection from Roundworms

ACTIVE INGREDIENT: 2.27 mg of pyrantel base as pyrantel pamoate per mL

NADA #200-248 • Approved by FDA
NET CONTENTS: 60 mL (2.11 oz)

DIRECTIONS FOR USE

For maximum control and prevention of reinfection, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfection. Administer one full teaspoonful (5 mL) for each 5 lb of body weight.

For the removal of large roundworms (ascarids) and hookworms. Administer one full teaspoonful (5 mL) for each 5 lb of body weight. It is not necessary to withhold food prior to treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infection should have a follow-up fecal exam within 2 to 4 weeks after first treatment.

Manufactured for:
Farnam Pet Products
 A Division of Farnam Companies, Inc.
 P.O. Box 34620 • Phoenix, AZ 85067-4620
 For information or comments, call 1-800-234-2269
 Visit our Web site at farnampet.com



Farnam Pet Products • A Division of Farnam Companies, Inc.
P.O. Box 34820 • Phoenix, AZ 85067-4820


D-Worm™ 60 *(pyrantel pamoate)*

Liquid Dewormer for Puppies & Dogs

ANADA #200-248 • Approved by FDA
FOR ANIMAL USE ONLY

D-Worm™ 60 (pyrantel pamoate suspension 2.27 mg) is a suspension of pyrantel pamoate in a palatable vanilla-flavored vehicle. Each mL contains 2.27 mg of pyrantel base as pyrantel pamoate.

Pyrantel pamoate is a compound belonging to a family classified chemically as tetrahydropyrimidines. It is a yellow, water-insoluble crystalline salt of the tetrahydropyrimidine base and pamoic acid containing 34.7% base activity. The chemical structure and name are given below:



(E)-1,4,5,6-Tetrahydro-1-methyl-2-[2-(2-thienyl) vinyl] pyrimidine 4,4' methylenebis [3-hydroxy-2-naphthoate](1:1)

INDICATIONS FOR USE

D-Worm™ 60 (*pyrantel pamoate*) suspension is a highly palatable formulation intended as a single treatment for the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies.

D-Worm™ 60 (*pyrantel pamoate*) suspension may also be used to prevent reinfection of *Toxocara canis* in puppies and adult dogs and in lactating bitches after whelping. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

DOSAGE AND ADMINISTRATION

Administer one full teaspoon (5 mL) for each 5 pounds of body weight (2.27 mg base per lb of body weight). Although most dogs have been observed to find this formulation very palatable and willingly consume it undiluted, it may be necessary to mix a small quantity of formulation in the dog's normal ration to encourage consumption. Fasting prior to or after treatment is not necessary.

For maximum control and prevention of reinfection, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfection.

CAUTION: THIS PRODUCT IS A SUSPENSION AND AS SUCH WILL SEPARATE. TO ENSURE UNIFORM RE-SUSPENSION AND TO ACHIEVE PROPER DOSAGE, **IT IS EXTREMELY IMPORTANT THAT THE PRODUCT BE SHAKEN THOROUGHLY BEFORE EVERY USE.**

RECOMMENDED STORAGE: Store below 86°F (30°C).

EFFICACY

Critical (worm count) studies in dogs demonstrated that pyrantel pamoate suspension 2.27 mg at the recommended dosage is highly efficacious against *T. leonina* (99%), *T. canis* (85%), *A. caninum* (97%), and *U. stenocephala* (94%).

SAFETY

One of the most outstanding and significant features of D-Worm™ 60 is its wide margin of therapeutic safety in dogs. The acute oral LD₅₀ of pyrantel pamoate administered in gelatin capsules to female and male dogs is greater than 314 mg base per pound of body weight, which indicates a therapeutic index in excess of 138 x the recommended dosage. In subacute and chronic studies, no significant morphological abnormalities could be attributed to pyrantel pamoate suspension 2.27 mg when administered to dogs at daily dose rates up to 94 mg base per pound of body weight (40 x) for periods of 19, 30, and 90 days. Clinical studies conducted in a wide variety of geographic locations using more than 40 different breeds of dogs showed no drug-induced toxic effects. Included in these studies were nursing pups, weaned pups, adults, pregnant bitches, and males at stud. Additional data have demonstrated the safe use of pyrantel pamoate in (1) dogs having heartworm infections and/or receiving medication for heartworm, (2) dogs exposed to organophosphate flea collars or flea/tick dip treatments, and (3) dogs undergoing concurrent treatment or medication at the time of worming such as immunization and antibacterial treatment.



07-1592

Distributed by:
Farnam Pet Products • A Division of Farnam Companies, Inc.
P.O. Box 34820 • Phoenix, AZ 85067-4820

500021

Rev0807





INDICATIONS FOR USE

To prevent reinfection of *Toxocara canis* in puppies and adult dogs and in lactating bitches after whelping. For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal examination. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

ACTIVE INGREDIENT: 4.54 mg of pyrantel base as pyrantel pamoate per mL.



Manufactured for:
Farnam Pet Products
A Division of Farnam Companies, Inc.
P.O. Box 34820
Phoenix, AZ 85067-4820
For information or comments, call
1-800-234-2269
Visit our Web site at farnampet.com

Product Code: 3087 500015 N-Rev0707
©2007 Farnam Companies, Inc. 07-1453

DOUBLE STRENGTH NDC 017135-677-12

D-Worm™ 120
(pyrantel pamoate)

Hookworms & Large Roundworms
Liquid
for Puppies & Dogs



Dog Dewormer

Easy-to-use

- Removes Hookworms and Large Roundworms (Ascarids)
- Prevents reinfection from Roundworms



5 ml per 10 lb Body Weight

ANADA #200-248 • Approved by FDA
NET CONTENTS: 60 mL (2 fl oz)



CAUTION:
IT IS EXTREMELY IMPORTANT
THAT THE PRODUCT BE SHAKEN THOROUGHLY
BEFORE EVERY USE.

DIRECTIONS FOR USE

For maximum control and prevention of reinfection, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfection. Administer one full teaspoonful (5 mL) for each 10 lb of body weight.

For the removal of large roundworms (ascarids) and hookworms. Administer one full teaspoonful (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to treatment. Dogs usually find this wormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption.

It is recommended that dogs maintained under conditions of constant exposure to worm infection should have a follow-up fecal exam within 2 to 4 weeks after first treatment.

**IMPORTANT: SHAKE WELL BEFORE USE
READ ACCOMPANYING PACKAGE INSERT**

**FOR ANIMAL USE ONLY
RECOMMENDED STORAGE:
Store Below 86° F (30° C).**



DOUBLE STRENGTH NDC 017135-677-12

D-Worm™ 120
(pyrantel pamoate)

Hookworms & Large Roundworms
Liquid
for Puppies & Dogs



Dog Dewormer

Easy-to-use

- Removes Hookworms and Large Roundworms (Ascarids)
- Prevents reinfection from Roundworms



5 ml per 10 lb Body Weight

ANADA #200-248 • Approved by FDA
NET CONTENTS: 60 mL (2 fl oz)



**Hookworms & Large Roundworms
Liquid Dewormer for Puppies & Dogs**

DOUBLE STRENGTH

5 ml per 10 lb Body Weight

**Hookworms & Large Roundworms
Liquid Dewormer for Puppies & Dogs**

DOUBLE STRENGTH

5 ml per 10 lb Body Weight

INDICATIONS FOR USE: To prevent reinfection of *Toxocara canis* in puppies and adult dogs and in lactating bitches after whelping. For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncia stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by bloody fecal examination. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasites.

IMPORTANT: SHAKE WELL BEFORE USE
READ ACCOMPANYING PACKAGE INSERT
FOR ANIMAL USE ONLY
RECOMMENDED STORAGE:
 Store below 86° F (30° C).

Product Code: 3087
 50015 Rev0807
 ©2007 Farnam Companies, Inc.
 07-1989

DOUBLE STRENGTH
D-Worm™ 120
(pyrantel pamoate)
Dog Dewormer
 Liquid for Puppies & Dogs

Easy-to-use:

- Removes Hookworms and Large Roundworms (Ascarids)
- Prevents reinfection from Roundworms

ACTIVE INGREDIENT: 4.54 mg of pyrantel base as pyrantel pamoate per mL

NET CONTENTS: 60 mL (2.11 oz)

MAJADA #200-248 • Approved by FDA

DIRECTIONS FOR USE

For maximum control and prevention of reinfection, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfection. Administer one full teaspoonful (5 mL) for each 10 lb of body weight.

For the removal of large roundworms (ascarids) and hookworms, Administer one full teaspoonful (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infection should have a follow-up fecal exam within 2 to 4 weeks after first treatment.

Manufactured for:
Farnam Pet Products
 A Division of Farnam Companies, Inc.
 P.O. Box 34620 • Phoenix, AZ 85067-4620
 For information or comments, call 1-800-234-2269
 Visit our Web site at farnampet.com


WARNING-KEEP OUT OF REACH OF CHILDREN



Farnam Pet Products • A Division of Farnam Companies, Inc.
P.O. Box 34820 • Phoenix, AZ 85067-4820

DOUBLE STRENGTH
D-Worm™ 120
(pyrantel pamoate) **Liquid Dewormer**
for Puppies & Dogs
ANADA #200-248 • Approved by FDA
FOR ANIMAL USE ONLY

D-Worm™ 120 (pyrantel pamoate suspension 4.54 mg) is a suspension of pyrantel pamoate in a palatable vanilla-flavored vehicle. Each mL contains 4.54 mg of pyrantel base as pyrantel pamoate. Pyrantel pamoate is a compound belonging to a family classified chemically as tetrahydropyrimidines. It is a yellow, water-insoluble crystalline salt of the tetrahydropyrimidine base and pamoic acid containing 34.7% base activity. The chemical structure and name are given below:



(E)-1,4,5,6-Tetrahydro-1-methyl-2-[2-(2-thienyl) vinyl] pyrimidine 4,4' methylenebis [3-hydroxy-2-naphthoate](1:1)

INDICATIONS FOR USE

D-Worm™ 120 (*pyrantel pamoate*) suspension is a highly palatable formulation intended as a single treatment for the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies.

D-Worm™ 120 (*pyrantel pamoate*) suspension may also be used to prevent reinfection of *Toxocara canis* in puppies and adult dogs and in lactating bitches after whelping. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

DOSAGE AND ADMINISTRATION

Administer one full teaspoon (5 mL) for each 10 pounds of body weight (2.27 mg base per lb of body weight). Although most dogs have been observed to find this formulation very palatable and willingly consume it undiluted, it may be necessary to mix a small quantity of formulation in the dog's normal ration to encourage consumption. Fasting prior to or after treatment is not necessary.

For maximum control and prevention of reinfection, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfection.

CAUTION: THIS PRODUCT IS A SUSPENSION AND AS SUCH WILL SEPARATE. TO INSURE UNIFORM RE-SUSPENSION AND TO ACHIEVE PROPER DOSAGE, **IT IS EXTREMELY IMPORTANT THAT THE PRODUCT BE SHAKEN THOROUGHLY BEFORE EVERY USE.**

RECOMMENDED STORAGE: Store below 86°F (30°C).

EFFICACY

Critical (worm count) studies in dogs demonstrated that pyrantel pamoate suspension 4.54 mg at the recommended dosage is highly efficacious against *T. leonina* (99%), *T. canis* (85%), *A. caninum* (97%), and *U. stenocephala* (94%).

SAFETY

One of the most outstanding and significant features of D-Worm™ 120 is its wide margin of therapeutic safety in dogs. The acute oral LD₅₀ of pyrantel pamoate administered in gelatin capsules to female and male dogs is greater than 314 mg base per pound of body weight, which indicates a therapeutic index in excess of 138 x the recommended dosage. In subacute and chronic studies, no significant morphological abnormalities could be attributed to pyrantel pamoate suspension 4.54 mg when administered to dogs at daily dose rates up to 94 mg base per pound of body weight (40 x) for periods of 19, 30, and 90 days. Clinical studies conducted in a wide variety of geographic locations using more than 40 different breeds of dogs showed no drug-induced toxic effects. Included in these studies were nursing pups, weaned pups, adults, pregnant bitches, and males at stud. Additional data have demonstrated the safe use of pyrantel pamoate in (1) dogs having heartworm infections and/or receiving medication for heartworm, (2) dogs exposed to organophosphate flea collars or flea/tick dip treatments, and (3) dogs undergoing concurrent treatment or medication at the time of worming such as immunization and antibacterial treatment.



07-1591

Distributed by:
Farnam Pet Products • A Division of Farnam Companies, Inc.
P.O. Box 34820 • Phoenix, AZ 85067-4820

500015

Rev0807

